

cond-mat/9406056

UCSB-TH-94-17

June 1994

Quantum Chaos and Statistical Mechanics

MARK SREDNICKI

Department of Physics

University of California

Santa Barbara, CA 93106

`mark@tpau.physics.ucsb.edu`

ABSTRACT: We briefly review the well known connection between classical chaos and classical statistical mechanics, and the recently discovered connection between quantum chaos and quantum statistical mechanics.

*Talk given at the Conference on
Fundamental Problems in Quantum Theory
Baltimore, June 18-22, 1994*

Consider a dilute gas of hard spheres in a box with hard walls. Give the spheres some arbitrary initial distribution of momenta (and positions). Classically, after a few mean free times have passed, we expect that the distribution of momenta will be given by the Maxwell–Boltzmann (MB) formula,

$$f_{\text{MB}}(\mathbf{p}) = (2\pi mkT)^{-3/2} \exp(-\mathbf{p}^2/2mkT) , \quad (1)$$

where the temperature T is given in terms of the conserved total energy U by the ideal-gas relation $U = \frac{3}{2}NkT$.

To see why this should be so, first note that the hamiltonian is simply

$$H = \frac{1}{2m} \sum_{i=1}^N \mathbf{p}_i^2 = \frac{1}{2m} \mathbf{P}^2 , \quad (2)$$

where $\mathbf{P}$ is a vector with $3N$ components. Since H takes on the constant value U , the allowed values of $\mathbf{P}$ form a sphere which we will call the P-sphere. Suppose we now choose $\mathbf{P}$ “at random.” For this to be a meaningful statement, we must have a measure which tells us which sets of $\mathbf{P}$ ’s are equally likely *a priori*. The obvious choice is to assign equal *a priori* probabilities to equal areas on the P-sphere. Then if we choose $\mathbf{P}$ at random with respect to this measure, the probability that our choice makes an angle between θ and $\theta + d\theta$ with respect to any particular axis is simply

$$\begin{aligned} f(\theta) d\theta &\sim (\sin \theta)^{3N-2} d\theta \\ &\sim (\sin \theta)^{3N-3} d\cos \theta \\ &\sim (1 - \cos^2 \theta)^{(3N-3)/2} d\cos \theta . \end{aligned} \quad (3)$$

If we now identify $(2mU)^{1/2} \cos \theta$ as, say, the value of p_{1z} (the z component of the first particle’s momentum), we find

$$\begin{aligned} f(p_{1z}) dp_{1z} &\sim (1 - p_{1z}^2/2mU)^{(3N-3)/2} dp_{1z} \\ &\sim \exp(-p_{1z}^2/2mkT) dp_{1z} , \end{aligned} \quad (4)$$

where in the second line we have set $U = \frac{3}{2}NkT$ and taken the large- N limit. Thus we have recovered the MB distribution for p_{1z} . Now consider the probability distribution for p_{1y} when p_{1z} is fixed; it is given by the first line of (4) with $3N$ replaced by $3N - 1$ (since there is one less coordinate when p_{1z} is fixed) and $2mU$ replaced by $2mU - p_{1z}^2$. In the large- N limit, we can neglect p_{1z}^2 compared to $2mU$, and we find the MB distribution for

p_{1y} . In similar fashion, we get the MB distribution for any n components of $\mathbf{P}$ as long as $n \ll N$.

Now our task is to justify the assumption that equal areas on the P-sphere are equally likely *a priori*. That such a justification is needed can be seen by considering how we would go about filling a real box with a real gas (say, helium). If the box did not already have some sort of valve on it, we would install one, and pump the air out through it. Then we would close the valve, attach it to a tank of helium with a hose, and open the valve. The helium atoms would rush in, moving preferentially in the direction parallel to the hose. Thus their initial distribution of momenta would be strongly anisotropic. This is in sharp contrast to the prediction of the equal-area measure, which tells us that we will find a thermal, isotropic distribution. Clearly, then, the equal-area measure has nothing to do with how we put real gases in real boxes, and so we must seek its justification elsewhere.

That justification comes from Sinai's theorem¹, which states that a box of hard spheres is a *chaotic* system. The meaning of this statement in the present context is simple. Start off with arbitrary initial momenta and positions; the momenta can be as nonthermal as you like. (Actually, we must exclude a set of measure zero; for example, it is possible to set up initial conditions such that no two hard spheres ever collide, in which case the following discussion obviously does not apply.) Wait a few mean free times, and then note the current location of $\mathbf{P}$ on the P-sphere. Continue this procedure, keeping track of the location of $\mathbf{P}$ each time. Chaos implies that this sequence of $\mathbf{P}$'s appears to be chosen at random with respect to the equal-area measure.

We are done. Even if we started off with a $\mathbf{P}$ representing a strongly anisotropic distribution, the next $\mathbf{P}$ will appear to be chosen "at random," and so predicts a thermal distribution for the individual momenta.

So much for classical mechanics. What about quantum mechanics?

Now we have a completely different problem². The N -particle Schrödinger equation can always be solved by going to the energy eigenstate basis: $H|\alpha\rangle = U_\alpha|\alpha\rangle$. The hamiltonian is given by (2), supplemented by the boundary condition that the energy eigenfunctions $\psi_\alpha(\mathbf{X})$ vanish whenever one of the hard spheres touches a wall of the box, or whenever two hard spheres touch each other. The wave function in momentum space at time t is then

$$\tilde{\psi}(\mathbf{P}, t) = \sum_{\alpha} C_{\alpha} \exp(-iU_{\alpha}t/\hbar) \tilde{\psi}_{\alpha}(\mathbf{P}) , \quad (5)$$

where the C_{α} 's specify the initial state. The probability that the first particle has momentum $\mathbf{p}_1$ at time t is found by squaring the wave function and integrating over all momenta

but the first:

$$\begin{aligned}
f(\mathbf{p}_1, t) &= \int d^3 p_2 \dots d^3 p_N |\tilde{\psi}(\mathbf{P}, t)|^2 \\
&= \sum_{\alpha\beta} C_\alpha^* C_\beta e^{i(U_\alpha - U_\beta)t/\hbar} \int d^3 p_2 \dots d^3 p_N \tilde{\psi}_\alpha^*(\mathbf{P}) \tilde{\psi}_\beta(\mathbf{P}) \\
&= \sum_{\alpha\beta} C_\alpha^* C_\beta e^{i(U_\alpha - U_\beta)t/\hbar} \Phi_{\alpha\beta}(\mathbf{p}_1) .
\end{aligned} \tag{6}$$

In the last line we have introduced

$$\Phi_{\alpha\beta}(\mathbf{p}_1) \equiv \int d^3 p_2 \dots d^3 p_N \tilde{\psi}_\alpha^*(\mathbf{P}) \tilde{\psi}_\beta(\mathbf{P}) , \tag{7}$$

which obeys the normalization condition

$$\int d^3 p_1 \Phi_{\alpha\beta}(\mathbf{p}_1) = \delta_{\alpha\beta} . \tag{8}$$

If we symmetrize or antisymmetrize each $\tilde{\psi}_\alpha(\mathbf{P})$ on exchange of any two $\mathbf{p}_i$'s to reflect Bose–Einstein (BE) or Fermi–Dirac (FD) statistics, then $f(\mathbf{p}_1, t)$ is independent of which $\mathbf{p}_i$ we choose.

On physical grounds, we expect that $f(\mathbf{p}_1, t)$ should be the MB (or BE or FD) distribution for any time t greater than a few mean free times. It is not obvious how this can occur. Consider, for example, what happens if we take the infinite time average of $f(\mathbf{p}_1, t)$:

$$\lim_{\tau \rightarrow \infty} \frac{1}{\tau} \int_0^\tau dt f(\mathbf{p}_1, t) = \sum_\alpha |C_\alpha|^2 \Phi_{\alpha\alpha}(\mathbf{p}_1) . \tag{9}$$

The infinite time average is obviously not something we can actually observe, but theoretically, if anything is going to be thermal, this is it. The problem is that the C_α 's are essentially arbitrary, so how can we possibly get the MB distribution?

There is only one way: each $\Phi_{\alpha\alpha}(\mathbf{p}_1)$ must *individually* be equal to the MB (or BE or FD) distribution at a temperature T_α which is given (at least approximately) by the ideal-gas relation $U_\alpha = \frac{3}{2} N k T_\alpha$. We call this hypothesis *eigenstate thermalization*. If eigenstate thermalization is valid, then (9) will indeed be a thermal distribution as long as the uncertainty in the total energy is much less than its expectation value.

Furthermore, if $\Phi_{\alpha\beta}(\mathbf{p}_1)$ is always sufficiently small whenever $\alpha \neq \beta$, then the $\alpha \neq \beta$ terms in (6) will usually make a negligible contribution, and $f(\mathbf{p}_1, t)$ will be a thermal distribution at most times t , without any time averaging at all. However, if the magnitudes

and phases of the C_α 's are carefully chosen, then we can “line up” the $\Phi_{\alpha\beta}(\mathbf{p}_1)$'s so as to get any $f(\mathbf{p}_1, t)$ that we might want at any one particular time (say, $t = 0$). Afterward, however, as we see in (6), the phases will change in the usual manner; the carefully contrived coherence among the various $\Phi_{\alpha\beta}(\mathbf{p}_1)$'s will be destroyed, and we will again find a thermal distribution for $\mathbf{p}_1$.

I find this to be a clear and satisfying explanation for the validity of quantum statistical mechanics, at least in this particular problem, even without any further evidence in favor of it. However, there is more to be said: a very strong case can be made for the two necessary ingredients—the thermal nature of $\Phi_{\alpha\alpha}(\mathbf{p}_1)$ and the smallness of $\Phi_{\alpha\beta}(\mathbf{p}_1)$ —based on the theory of quantum chaos.

Quantum chaos is the study of quantum systems whose classical counterparts are chaotic. The result we will need is known as Berry's conjecture^{3–5}. As its name implies, Berry's conjecture is as yet unproved, but there is significant numerical evidence (reviewed in [2]) in support of it.

Berry's conjecture has two parts. Part one says that the energy eigenfunctions of a bounded, isolated quantum system which is classically chaotic appear to be gaussian random variables, in the sense that

$$\lim_{\alpha \rightarrow \infty} \int d\mathbf{X} \, \psi_\alpha(\mathbf{X} + \mathbf{X}_1) \dots \psi_\alpha(\mathbf{X} + \mathbf{X}_n) = \sum_{\text{pairs}} J(\mathbf{X}_{i_1} - \mathbf{X}_{i_2}) \dots J(\mathbf{X}_{i_{n-1}} - \mathbf{X}_{i_n}) . \quad (10)$$

Here the integration measure is normalized so that $\int d\mathbf{X} = 1$, and the sum is over all possible ways to pair up the $\mathbf{X}_i$'s; if n is odd the result is zero. Part two says that the correlation function $J(\mathbf{X})$ is given by

$$J(\mathbf{X}) \sim \int d\mathbf{P} \, \exp(i\mathbf{P} \cdot \mathbf{X} / \hbar) \, \delta(H(\mathbf{P}, \mathbf{X}) - U_\alpha) , \quad (11)$$

where $\delta(x)$ is the Dirac delta function, and $J(\mathbf{0}) = 1$.

It is straightforward to show that Berry's conjecture gives us the two necessary ingredients for quantum statistical mechanics. First, we find that when $\alpha \neq \beta$, $\Phi_{\alpha\beta}(\mathbf{p}_1)$ is exponentially small in the number of particles N . More importantly, we find eigenstate thermalization: $\Phi_{\alpha\alpha}(\mathbf{p}_1)$ is given by the MB or BE or FD distribution (depending on whether we use nonsymmetric, completely symmetric, or completely antisymmetric energy eigenfunctions), plus corrections which depend on the specific energy eigenfunction but which are exponentially small in N . To derive this, the gas must be dilute; there are also other, hard-to-compute corrections to $\Phi_{\alpha\alpha}(\mathbf{p}_1)$ due to the finite radii of the hard spheres. We expect these to reproduce the usual hard-sphere corrections to ideal-gas behavior, but this remains to be demonstrated. Another important unsettled issue is

how high the energy needs to be before (10) is sufficiently accurate. A naive estimate is $\lambda_\alpha \lesssim a$, where $\lambda_\alpha = (2\pi\hbar^2/mkT_\alpha)^{1/2}$ is a typical particle's de Broglie wavelength, and a is the hard-sphere radius. With a in angstroms and m in amu, this condition becomes $T_\alpha \gtrsim (300/ma^2)$ Kelvin.

To summarize, the appearance of a thermal distribution of momenta in an isolated, bounded quantum system of many particles can only be understood if each energy eigenfunction individually predicts a thermal probability for the momentum of each constituent particle, and if overlaps of different energy eigenfunctions are sufficiently small when one particle's momentum is left unintegrated. Both these statements can be derived as consequences of Berry's conjecture, which is expected to hold only for quantum systems whose classical counterparts are chaotic. Thus the well known connection between classical statistical mechanics and classical chaos is now seen to be mirrored by an analogous connection between quantum statistical mechanics and quantum chaos.

I would like to thank Edouard Brezin, John Cardy, Marty Halpern, and Walter Kohn for helpful discussions. This work was supported in part by NSF Grant PHY-91-16964.

References

1. Ya. G. Sinai, *Ups. Mat. Nauk* 25, 137 (1970) [*Russ. Math. Surv.* 25, 137 (1970)].
2. M. Srednicki, cond-mat/9403051, *Phys. Rev. E*, in press.
3. M. V. Berry, *J. Phys. A* 10, 2083 (1977).
4. M. V. Berry, in *Les Houches XXXVI, Chaotic Behavior of Deterministic Systems*, G. Iooss, R. H. G. Helleman, and R. Stora, eds. (North-Holland, Amsterdam, 1983).
5. M. V. Berry, in *Les Houches LII, Chaos and Quantum Physics*, M.-J. Giannoni, A. Voros, and J. Zinn-Justin, eds. (North-Holland, Amsterdam, 1991).